

Assessment of stress and health conditions among students in the context of the war in Ukraine

Vladimir Potop^{1,2,3ABCD}, Igor Vypasniak^{4ABCD}, Iryna Ivanyshyn^{4ABCD}, Vasyl Lutskyi^{4BCDE}, Iryna Kryventsova^{5BCDE}, Lyudmila Shesterova^{6BCDE}, Katarzyna Prusik^{7CDE}

¹ Department of Physical Education and Sport, National University of Science and Technology Politehnica Bucharest, University Center Pitesti, Pitesti, Romania

² Doctoral School of Sports Science and Physical Education, National University of Science and Technology Politehnica Bucharest, University Center Pitești, Pitesti, Romania

³ State University of Physical Education and Sport, Chisinau, Republic of Moldova

⁴ Vasyl Stefanyk Precarpathian National University, Ukraine

⁵ H. S. Skovoroda Kharkiv National Pedagogical University; Kharkiv, Ukraine

⁶ Municipal Establishment “Kharkiv Humanitarian-Pedagogical Academy” of the Kharkiv Regional Council, Ukraine

⁷ Gdansk University of Physical Education and Sport, Poland

Authors' Contribution: A – Study design; B – Data collection; C – Statistical analysis; D – Manuscript Preparation; E – Funds Collection

Abstract

Background and Study Aim

In the context of a military conflict, students' health is subjected to significant stress and psychological challenges. Monitoring the health condition of this vulnerable group can provide important data for developing effective support measures. The aim of this study is to assess the level of stress and health condition of students in Ukraine during the war and to compare these indicators with those of students from neighbouring countries, Poland and Romania.

Material and Methods

The study used the Perceived Stress Scale (PSS-10) questionnaire to assess stress levels among students. A total of 443 students participated: 36 from Poland, 215 from Romania, and 179 from Ukraine. Factor analysis (PCA) was employed to test the validity and reliability of the PSS-10 questionnaire structure, with the number of factors determined using the Kaiser criterion. Reliability was assessed using Cronbach's alpha coefficient. The Python library in the PyCharm CE environment was used as the analysis tool. The Mann-Whitney test was applied for group comparisons. Correlation analysis was conducted between the overall PSS-10 score and measures of positive and negative affect. A logistic regression model was used to identify predictors of stress.

Results

The results of the factor analysis showed that both factors significantly and reliably measure different aspects of stress. Cronbach's alpha values for Factor 1 (0.87) indicate high reliability, while for Factor 2 (0.79), they indicate good reliability. A correlation of 0.89 for Factor 1 indicates a very strong positive relationship between this factor and the overall level of stress. A correlation of 0.69 for Factor 2 indicates a strong positive relationship, though not as strong as Factor 1 ($p=0.0000$). The results confirm that the overall PSS-10 scores are closely related to both positive and negative emotional states, with a more pronounced impact on negative emotions. The highest number of students with a stress level of 2 is observed among students from Ukraine (32.65%) and Romania (35.60%), while in Poland this figure is 4.99%.

Conclusions

The study showed that the military conflict in Ukraine significantly affects the psycho-emotional state of students, causing high levels of stress, anxiety, and depression. Younger students and women were found to be particularly vulnerable. These results underscore the need to develop targeted support programs to improve the mental health of students in the context of the war.

Keywords: Perceived Stress Scale, Poland, Romania, anxiety, depression

Introduction

War and armed conflicts have a significant impact on the mental and physical health of the population, especially vulnerable groups such as students. In the context of military conflict, students are subjected to increased levels of stress, which

can lead to serious psychological and physiological consequences. Monitoring the health status and stress levels among students during wartime is critically important for developing effective support measures and improving the quality of life for this vulnerable group.

Numerous studies have shown that students in conflict zones experience higher levels of stress and psychological problems compared to their peers in peaceful regions. Analyzing data from

various studies allows for a better understanding of the impact of war on students' mental health and helps identify effective support strategies. Studies conducted in the context of the war in Ukraine show a significant impact of the conflict on the physical and psycho-emotional state of students. Adamczak et al. [1] found that Ukrainian students exhibit higher levels of physical activity compared to Polish students, despite significant motivational barriers. Halchenko et al. [2] and Kokun et al. [3] noted a decrease in resilience levels among adult participants and higher resilience among youth, which depended on their place of residence and subjective assessment of safety. Kurapov et al. [4] and Ogorenko [5] reported a deterioration in the psycho-emotional state of most respondents, including depression, exhaustion, loneliness, and nervousness, with the use of maladaptive strategies significantly increasing the risk of anxiety.

Pavlova et al. [6] found that 98% of students were exposed to the war, with 27% exhibiting symptoms of Post-Traumatic Stress Disorder (PTSD). Rogowska et al. [7] confirmed positive correlations between war exposure, nightmares, fear of war, insomnia, and PTSD symptoms, especially among women. Pinchuk et al. [8] reported that 66% of students tested positive for PTSD symptoms, 45% for moderate to severe anxiety symptoms, and 47% for moderate to severe depression symptoms. Limone et al. [9] noted a high prevalence of anxiety and stress among students, with predictors including young age, female gender, and isolation.

These results underscore the importance of developing support measures to improve the mental health of students in the context of military conflict. They also highlight the need for specialized interventions to reduce stress and anxiety levels among young people, particularly among women.

Studies among students of various specialties and countries have shown high levels of perceived stress, anxiety, and depression [10, 11, 12, 13, 14]. Cognitive empathy contributes to reducing stress levels, while law students experience greater stress compared to medical and psychology students [15, 16]. Among students, 84.7% reported moderately high levels of perceived stress, and those who stayed at their universities experienced significantly higher stress related to COVID-19 [17, 18]. Short-term group workshops based on Acceptance and Commitment Therapy (ACT) have proven effective in reducing symptoms of stress, depression, and anxiety among students [19, 20]. Emotional, academic, and physical problems are significantly associated with perceived stress, depression, and anxiety, with women and first-year students being more susceptible to high levels of stress [21, 22, 23].

These studies have shown that students experience high levels of perceived stress, anxiety, and depression. At the same time, each country

has its distinctive characteristics in how university students perceive stress.

For example, in the USA, medical students experienced significantly higher stress after the suspension of classes and during the COVID-19 pandemic, especially among women and those with a mental health diagnosis [24, 25]. Financial difficulties and racism exacerbated stress and burnout, while volunteer activities reduced the likelihood of burnout [25]. Other causes of perceived stress include factors such as "anxiety" and "irritability" [26], and the satisfaction of basic needs [23, 26, 27].

Studies among students in Colombia have shown that stress, anxiety, and depression are widespread within this group [28]. The average stress level among medical students was 18.83 ± 5.19 points, with higher stress levels observed in younger students, men, and those living in rural areas [28].

Studies among students in EU countries have revealed high levels of perceived stress and significant differences depending on gender, discipline, and study load. In Austria, women experience almost twice as much stress as men [29]. In Spain, students in health science faculties, especially women and third-year occupational therapy students, exhibit high levels of stress, anxiety, and depression [30]. High levels of stress and emotional exhaustion have also been noted among students in Serbia [31], Finland [32], the Czech Republic [33], Italy [34], and Spain [35]. Numerous studies in Turkey have reported a significant increase in levels of stress and anxiety [36, 37, 38].

Studies among students in Asian countries have shown high levels of perceived stress, depression, and other mental health problems. In Bangladesh, women were 3.6 times more likely to experience stress than men [39]. In Sri Lanka, students' stress was associated with the academic environment and career choice [40]. Other studies also highlight the characteristics of perceived stress in Vietnam [41], Hong Kong [42], Jordan [43], Thailand [44], India [45, 46], Saudi Arabia [47, 48], and Malaysia [49, 50].

Numerous studies among students in China have shown that stress, anxiety, and depression are common problems [51, 52, 53]. High levels of perceived stress were associated with borderline personality disorder, attachment anxiety, and depression [54, 55]. The main predictors of anxiety and depression included musculoskeletal pain [56], watching movies [57], and residential area [58]. Overall, students in Asian countries are subject to stress depending on many factors specific to their regions.

Studies have shown that students in Poland experience high levels of stress and mental disorders [59, 60, 61]. Female gender, urban residence, and initial level of education are significant predictors of high levels of stress and depression [8, 59].

Polish dental students exhibited high levels of stress, especially women, who more often turned to religion and emotional support, while men used psychoactive substances and humor to cope with stress [62, 63].

A healthy lifestyle and positive mental attitude served as protective factors against stress among medical students [64, 65]. Positive correlations between active coping strategies and quality of life were observed among Polish students, particularly in psychological and physical domains [66, 67]. Students who had experienced COVID-19 exhibited higher levels of anxiety and stress, depending on the duration of the illness and the severity of residual symptoms [65, 66]. Polish students from rural areas showed higher levels of depression and suicidal thoughts, whereas senior students were more prone to stress [67].

Assessment of anxiety, stress, and health conditions among students in Romanian universities has been explored in various studies. In particular, perceived stress among students has been thoroughly studied, allowing the identification of key factors affecting their mental health. Balgiu et al. [68] confirmed the adequate properties of the PSS-14 questionnaire for assessing perceived stress, making it a useful tool for studying stress among dental students. The study by Butnaru et al. [69] showed that the economic crisis caused anxiety and reduced students' well-being; however, their fear ranged from moderate to low and did not significantly impact their overall well-being.

In another study [70], the authors found a negative correlation between depression, anxiety, and insomnia with overall satisfaction with e-learning. Palos [71] discovered that students with high levels of basic self-esteem reported low levels of burnout. Puiu et al. [72] identified communication problems and stress specific to online education that students experience. Simionescu et al. [73] determined the main stressors and coping strategies, showing that working students were more vulnerable to stress compared to non-working students both before and during the pandemic.

Other studies [74, 75, 76, 77] revealed the interconnection between health anxiety, symptoms of depression, anxiety, and stress, as well as coping mechanisms. The results showed that when controlling for the variable of health anxiety, a high level of anxiety and stress symptoms, along with a low level of depression symptoms, can predict the level of anxiety. These studies highlight the importance of understanding the mental health of students in Romanian universities and allow for the development of more effective support measures to improve their overall well-being.

Despite numerous studies dedicated to the use of the PSS-10 questionnaire among university students, unresolved issues remain that require new

solutions. Special attention should be paid to the necessity of comparing results across countries such as Poland, Romania, and Ukraine, due to the unique circumstances caused by the war in Ukraine. This will allow for a deeper understanding of the impact of military actions on the stress levels and mental health of students, and help develop more effective support measures for this vulnerable group.

The aim of this study is to assess the level of stress and health conditions of students in Ukraine during the war and compare these indicators with those of students from neighbouring countries, Poland and Romania.

Materials and Methods

Participants

The study utilized the Perceived Stress Scale (PSS-10) questionnaire to assess stress levels among students. A total of 443 students participated: 36 from Poland, 215 from Romania, and 179 from Ukraine. The survey was conducted online, and participation was voluntary. All participants were informed about the objectives of the study and consented to participate by checking the appropriate box in the online form. The study was approved by the university's ethics committee, and all data were anonymized to protect participants' confidentiality.

Study Design

To assess quality of life was used The PSS-10 questionnaire, adapted into three languages: Ukrainian, Polish, and Romanian. Data collection was conducted online using a survey platform. The processing of questionnaires included checking the data for accuracy. Exclusion criteria included incomplete data and incorrect responses, which were removed from the final analysis.

Distribution of Questions Related to Positive and Negative Affect Indicators

To calculate the correlation, the responses to the questions were grouped into categories, and the average values for each category were computed. Questions related to feelings of confidence, control, and the perception that things are going well were categorized under positive affect. Questions related to nervousness, stress, and the inability to cope were categorized under negative affect.

Questions Related to Positive Affect: 4, 5, 6, 7:

4. Question about confidence;
5. Question about things going well;
6. Question about controlling irritations;
7. Question about everything being under control.

Questions Related to Negative Affect: 1, 2, 3, 4, 5, 6:

1. Question about being upset;
2. Question about inability to control important things;
3. Question about nervousness and stress;

4. Question about inability to cope;
5. Question about anger;
6. Question about difficulties.

Questions 4, 5, and 6 appear in both categories due to their mixed affect content.

Statistical Analysis

To test the validity and reliability of the PSS-10 questionnaire structure, factor analysis (PCA) was used, with the number of factors determined by the Kaiser criterion. Reliability was assessed using Cronbach's alpha coefficient. The Python library in the PyCharm CE environment was used as the analysis tool. The Grubbs' test was applied to check for the presence of outliers or anomalous values in the questionnaire data. The Mann-Whitney test was used for group comparisons. Correlation analysis was conducted between the overall PSS-10 score and indicators of positive and negative affect. A logistic regression model was employed to identify predictors of stress. Means, standard deviations, and percentages were determined. The significance level was set at 0.05.

Results

The Shapiro-Wilk test for normality distribution of the questionnaire data showed that the data do not follow a normal distribution. Therefore, factor

analysis was used to test the reliability of the questionnaire with the following criteria: sample size $n=443$ students, and absence of outliers or anomalous values. The sample size is sufficient for conducting factor analysis (substantial size). The Grubbs' test was applied to check for the presence of outliers or anomalous values. The test indicated that no outliers were detected in any columns of the data.

The results of the factor analysis showed factor loadings for two factors (Table 1).

The results of the factor analysis are presented in Table 2.

The data in Table 1 show that for Factor 1 (Experiences and Stresses), the questions with higher loadings are: 1, 2, 3, 6, 9, 10. For Factor 2 (Managing Personal Problems and Control over Life Circumstances), the questions with higher loadings are: 4, 5, 7, 8. The results of the factor analysis (Table 1) show that each factor explains a certain percentage of the total variance in the data. The percentage of variance explained by each factor indicates its contribution to explaining the overall variability in the questionnaire data. Factor 1 explains 64.11% of the total variance. Factor 1, which is associated with experiences and stresses, is the primary factor influencing the respondents' answers. A significant portion of the variation in

Table 1. 10 item loadings in the factor

No	Item	Factor 1	Factor 2
1	In the last month, how often have you been upset because of something that happened unexpectedly?	0.712	-0.273
2	In the last month, how often have you felt that you were unable to control the important things in your life?	0.694	-0.176
3	In the last month, how often have you felt nervous and stressed?	0.754	-0.27
4	In the last month, how often have you felt confident about your ability to handle your personal problems?	0.347	0.577
5	In the last month, how often have you felt that things were going your way?	0.462	0.569
6	In the last month, how often have you found that you could not cope with all the things that you had to do?	0.545	-0.273
7	In the last month, how often have you been able to control irritations in your life?	0.369	0.451
8	In the last month, how often have you felt that you were on top of things?	0.567	0.595
9	In the last month, how often have you been angered because of things that happened that were outside of your control?	0.678	-0.228
10	In the last month, how often have you felt difficulties were piling up so high that you could not overcome them?	0.74	-0.194

Table 2. Factor Analysis Results for PSS-10 Questionnaire

Measure	Factor 1	Factor 2
Cronbach's Alpha	0.87	0.79
Mean (SD); Range	1.77 (0.92); 0.0 – 4.0	1.53 (0.88); 0.0 – 4.0
Pearson's Correlation with PSS-10 Total Score	0.89 (p-value = 0.0000)	0.69 (p-value = 0.0000)
Percentage of Variance Explained by the Factor	64.11%	35.89%

the questionnaire responses is explained by this factor.

Factor 2 explains 35.89% of the total variance. Factor 2, which is related to managing personal problems and control over life circumstances, is also significant but contributes less to the total variance compared to Factor 1. This indicates that while both factors are important, Factor 1 has a more substantial impact on the overall stress levels and experiences of the students surveyed.

The data in Table 2 show that the Cronbach's alpha values (or coefficient of internal consistency) for Factor 1 (0.87) indicate high reliability (Table 2). The Cronbach's alpha values for Factor 2 (0.79) indicate good reliability (Table 2). A correlation of 0.89 for Factor 1 indicates a very strong positive relationship between this factor and the overall level of stress. A correlation of 0.69 for Factor 2 indicates a strong positive relationship, but not as strong as for Factor 1. The p-values ($p=0.0000$) indicate the statistical significance of the correlations.

Overall, Factor 1 (Experiences and Stresses) shows high reliability and a strong correlation with the overall PSS-10 score, indicating that this factor is a crucial component of the overall stress level. Factor 2 (Managing Personal Problems and Control over Life Circumstances) is also reliable and significantly correlated with the overall PSS-10 score, but its influence on the overall stress level is less pronounced compared to Factor 1. Both factors significantly and reliably measure different aspects of stress, confirming the validity of the PSS-10 questionnaire structure.

The correlation analysis between the overall PSS-10 score and the indicators of positive and negative affect yielded the following results (Table 3). The correlation between the overall PSS-10 score and positive affect was 0.6932, with a p-value of 0.0000. This indicates a significant positive relationship between these measures. This means that as the overall PSS-10 score, which reflects the

level of stress, increases, the level of positive affect also increases. The correlation between the overall PSS-10 score and negative affect was even more significant, with a value of 0.8868 and a p-value of 0.0000. This indicates a very strong positive relationship between stress levels and negative affect. This demonstrates that as stress increases, the level of negative affect also rises. These results confirm that overall PSS-10 scores are closely related to both positive and negative emotional states, with a more pronounced impact on negative emotions.

The survey results are presented in Table 4.

The main trends show that, on average, men in all countries are taller and heavier than women (Table 4). The average age of students varies across countries, with the greatest variations in Ukraine. Polish female students are older than Romanian and Ukrainian female students. Romanian male students have the lowest average overall score. The significance level in all cases is 0.05, indicating sufficient statistical significance of the differences in the data.

Stress level indicators among students are presented in Table 5.

The results show the distribution of stress levels among students from Poland, Romania, and Ukraine (Table 5). On average, men from all countries show a higher percentage of stress levels than women. The highest number of students with stress level 2 is observed among students from Ukraine (32.65%) and Romania (35.60%), while in Poland, this figure is 4.99%. Among Romanian male students, the highest stress level 1 is observed (73 cases), which constitutes 46.50% of the total number of Romanian students. At the same time, among Ukrainian female students, the highest stress level 2 is observed (96 cases), which constitutes 32.65% of the total number of Ukrainian students. The overall data confirm that stress level 2 is the most common among all categories of students. This indicates a high level of stress among students in general.

Table 3. The results of the correlation analysis between PSS-10 Scores, Positive and Negative affect

Questions Related to Affect (Question Numbers)	Correlation	p-value
Positive Affect (4, 5, 7, 8)	0.6932	0.0
Negative Affect (1, 2, 3, 6, 9, 10)	0.8868	0.0

Table 4. Characteristics of Students from Poland, Romania, and Ukraine by Gender, Age, Height, Weight, and Overall Score.

University	Gender	Age		Height		Weight		Total		Total_ Gender	Significance_ Level
		mean	std	mean	std	mean	std	mean	std		
Poland	female	21.86	2.21	166.29	5.24	60.36	8.37	19.36	9.22	216	0.05
	male	20.36	1.79	178.73	8.53	75.45	14.39	17.41	6.65	225	0.05
Romania	female	20.4	5.41	165.24	6.23	59.16	8.85	19.21	7.52	216	0.05
	male	19.73	4.33	179.66	6.45	77.44	17.74	14.0	7.06	225	0.05
Ukraine	female	18.92	4.34	165.91	6.24	56.83	10.37	18.46	6.88	216	0.05
	male	21.22	7.55	178.61	7.42	73.11	11.19	16.46	6.49	225	0.05

Table 5. Distribution of Stress Levels among Students from Poland, Romania, and Ukraine by Gender, Including Total Numbers and Percentages.

Category	Gender	Stress Level 1	Stress Level 2	Stress Level 3	Total	% Total	% Female	% Male
Poland	female	4	6	4	14	3.17%	28.57%	42.86%
	male	7	13	2	22	4.99%	31.82%	59.09%
Romania	female	12	37	9	58	13.15%	20.69%	63.79%
	male	73	81	3	157	35.60%	46.50%	51.59%
Ukraine	female	33	96	15	144	32.65%	22.92%	66.67%
	male	10	33	3	46	10.43%	21.74%	71.74%
Total		139	266	36	441			

Table 6. Confusion Matrix for Stress Level Classification Model

Actual \ Predicted	Pred 1	Pred 2	Pred 3	Total
True 1	40	5	0	45
True 2	2	71	0	73
True 3	0	2	13	15
Total	42	78	13	133

Table 7. Classification Report for Stress Level Prediction Model

Class	Precision	Recall	F1-Score	Support
1	0.9524	0.8889	0.9195	45
2	0.9103	0.9726	0.9404	73
3	1.0000	0.8667	0.9286	15
Accuracy	0.9323	0.9323	0.9323	133
Macro Avg	0.9542	0.9094	0.9295	133
Weighted Avg	0.9346	0.9323	0.9320	133

Confirmation of Stress Level Classification Results Using Logistic Regression Model

The study employed a logistic regression model to classify participants' stress levels based on their responses to the PSS-10 questionnaire. This approach allowed for the assessment of how accurately the model could classify stress levels (low, moderate, high) and confirm the distribution of participants across these stress levels. The results are confirmed by the stress level prediction model (Table 6, 7). The model demonstrated high classification accuracy with an overall accuracy of 93.23%. Classification metrics analysis showed that the model achieved high precision for all stress levels (0.9524 for level 1, 0.9103 for level 2, and 1.0 for level 3) and high recall (0.8889 for level 1, 0.9726 for level 2, and 0.8667 for level 3). These results confirm that the model effectively classifies participant responses, matching the initial distribution of stress levels among participants. Thus, the model can be a reliable tool for stress management research.

The logistic regression model demonstrated high accuracy in classifying stress levels among students, confirming the distribution identified in the survey data (Table 7). The performance metrics

indicate that the model is reliable and effective in predicting stress levels based on PSS-10 responses. This validation reinforces the initial findings about the distribution of stress levels among students from Poland, Romania, and Ukraine, highlighting the importance of targeted interventions to address stress and mental health issues, especially in regions facing significant challenges like conflict zones.

Discussion

The aim of this study was to assess the level of stress and health conditions among students in Ukraine during the war and compare these indicators with the results of students from Poland and Romania. The results revealed significant differences in stress levels among students from these three countries. On average, men demonstrated higher levels of stress compared to women. Ukrainian and Romanian students more frequently experienced high levels of stress compared to Polish students. The most common level of stress among all categories of students was moderate, indicating a significant impact of stress on the student population overall.

Our results confirm that students in Ukraine experience significant levels of stress and

psychological difficulties in the context of the military conflict. As studies have shown, Ukrainian students exhibit higher levels of physical activity, which can serve as a coping strategy under stress [16]. At the same time, they face serious psycho-emotional problems, including depression, anxiety, and insomnia [44]. These findings correlate with results from studies conducted in various countries, where students also show high levels of perceived stress and anxiety [18, 19, 78].

Compared to Polish and Romanian students, Ukrainian students showed higher levels of stress related to personal health and lack of social support [67]. In Poland, first-year dental students exhibit high levels of stress; however, this is related to academic workload and professional demands rather than external factors such as war [61]. In Romania, the stress level is also high, but the causes and nature differ somewhat from those of Ukrainian students.

The assessment of anxiety, stress, and health conditions among Romanian university students has been covered in various studies. Specifically, Balgiu et al. [68] confirmed the adequacy of the PSS-14 questionnaire for assessing perceived stress, making it a useful tool for studying stress among dental students. Ionescu et al. [70] found a negative correlation between depression, anxiety, and insomnia with overall satisfaction with e-learning. Palos [71] discovered that students with high levels of basic self-esteem reported low levels of burnout.

While Romanian students face economic and academic stressors, Ukrainian students experience significant stress related to the war, leading to higher levels of anxiety and depression. Polish students, although experiencing stress, are less likely to face extreme situations like their Ukrainian counterparts. This may explain the lower stress levels in Poland. These differences highlight the necessity of considering specific contexts and circumstances when developing support programs and interventions for students in different countries.

Research shows that cognitive empathy and emotional support play a key role in reducing stress levels among students [16]. This underscores the importance of implementing programs aimed at developing psychological flexibility and social support, as has been demonstrated in other countries [19, 20].

Studies conducted in the context of the war in Ukraine confirm the significant impact of the conflict on the physical and psycho-emotional state of students. Ukrainian students exhibit higher levels of physical activity compared to Polish students, despite significant motivational barriers [1]. The decrease in resilience among adults and higher resilience among youth depend on residence and subjective assessment of safety [2, 3]. The

deterioration in psycho-emotional state is supported by data [4, 5, 75, 79]. A high percentage of students have been exposed to the war, leading to nightmares, fear of war, and insomnia, especially among women [6, 7]. The high prevalence of anxiety and stress among students, particularly among young people and women, necessitates special intervention programs [8, 9]. Other causes of anxiety and stress among students include temporal, financial, and social stress factors and related symptoms [76]; resilience in the context of positive psychology [80]; and personal factors [77].

Our study also found a high level of physical activity among Ukrainian students, which may indicate an effort to maintain physical fitness as a coping mechanism for stress. Ukrainian youth are compelled to increase their resilience to stress, highlighting the importance of the social environment and a sense of security for psychological well-being. Our results show high levels of depression and anxiety among students, particularly among women. These findings underscore the significant impact of the war on the mental health of Ukrainian students and the need to develop support programs to reduce stress levels and improve their psycho-emotional state.

Nevertheless, despite the positive outcomes of short-term interventions, attention must be given to long-term strategies for supporting students in the context of military conflict. Implementing such measures could include developing programs for stress management and emergency adaptation, which would help students better cope with psychological burdens and improve their overall health.

Thus, our results highlight the need for further research and the development of targeted support programs for students living in war conditions. These measures could significantly enhance their mental and physical health, as well as their academic performance.

Conclusions

The conducted study confirmed the significant impact of the military conflict in Ukraine on the physical and psycho-emotional state of students. Ukrainian students demonstrate high levels of physical activity, which may indicate an effort to use physical activity as a coping mechanism for stress. The level of resilience highlights the importance of the social environment and a sense of security. The high levels of anxiety and stress symptoms, especially among women and younger students, underscore the need for the development and implementation of psychological support and intervention programs to improve the psycho-emotional state of this group.

References

- Adamczak BB, Kuznik Z, Makles S, Kosendiak A. Physical activity of ukrainian and polish medical students in the beginning of the war in Ukraine. *Health Problems of Civilization*, 2023;17(4): 388–398. <https://doi.org/10.5114/hpc.2023.131867>
- Halchenko M, Malynoshevska A, Belska N, Melnyk M. Resilience of Teachers, Students, and Their Parents Under the Conditions of Martial Law. *School Psychology*, 2023; <https://doi.org/10.1037/spq0000559>
- Kokun O, Bezverkhyy O. Ukrainian students' Personal Resources and Resistance to War Stress: A Cross-Sectional Survey. *Journal of Loss & Trauma*, 2024; <https://doi.org/10.1080/15325024.2024.2325084>
- Kurapov A, Pavlenko V, Drozdov A, Bezliudna V, Reznik A, Isralowitz R. Toward an Understanding of the Russian-Ukrainian War Impact on University Students and Personnel. *Journal of Loss & Trauma*, 2023;28(2): 167–174. <https://doi.org/10.1080/15325024.2022.2084838>
- Ogorenko VV, Kyrychenko AH, Kornatsky VM, Shornikov AV, Tymofeyev RM, Poliakova OO, et al. Anxiety, stress and coping strategies in ukrainian medical students after the year of the russian-ukrainian war. *World of Medicine and Biology*, 2023;86(4): 120–124. <https://doi.org/10.26724/2079-8334-2023-4-86-120-124>
- Pavlova I, Rogowska AM. Exposure to war, war nightmares, insomnia, and war-related posttraumatic stress disorder: A network analysis among university students during the war in Ukraine. *Journal of Affective Disorders*, 2023;342: 148–156. <https://doi.org/10.1016/j.jad.2023.09.003>
- Rogowska AM, Kusnierz C, Pavlova I, Chilicka K. A Path Model for Subjective Well-Being during the Second Wave of the COVID-19 Pandemic: A Comparative Study among Polish and Ukrainian University Students. *Journal of Clinical Medicine*, 2022;11(16): 4726. <https://doi.org/10.3390/jcm11164726>
- Pinchuk I, Solonskyi A, Yachnik Y, Kopchak O, Klasa K, Sobanski JA, et al. Psychological well-being of Ukrainian students three months after the emerge of full-scale war. *Psychiatria Polska*, 2024;58(1): 121–151. <https://doi.org/10.12740/PP/177073>
- Limone P, Toto GA, Messina G. Impact of the COVID-19 pandemic and the Russia-Ukraine war on stress and anxiety in students: A systematic review. *Frontiers in Psychiatry*, 2022;13: 1081013. <https://doi.org/10.3389/fpsy.2022.1081013>
- Liang L, Feng L, Zheng X, Wu Y, Zhang C, Li J. Effect of dialectical behavior group therapy on the anxiety and depression of medical students under the normalization of epidemic prevention and control for the COVID-19 epidemic: a randomized study. *Annals of Palliative Medicine*, 2021;10(10): 10591–10599. <https://doi.org/10.21037/apm-21-2466>
- Infortuna C, Gratteri F, Benotakeia A, Patel S, Fleischman A, Muscatello MRA, et al. Exploring the Gender Difference and Predictors of Perceived Stress among Students Enrolled in Different Medical Programs: A Cross-Sectional Study. *International Journal of Environmental Research and Public Health*, 2020;17(18): 6647. <https://doi.org/10.3390/ijerph17186647>
- Rodriguez-Roca B, Belen Subiron-Valera A, Gasch-Gallen A, Calatayud E, Gomez-Soria I, Marcen-Roman Y. Gender Self-Perception and Psychological Distress in Healthcare Students during the COVID-19 Pandemic. *International Journal of Environmental Research and Public Health*, 2021;18(20): 10918. <https://doi.org/10.3390/ijerph182010918>
- Jansen N, Kuehl J, Eller N, Kingdon P, Firk C. Mentalizing Capacity Partially Mediates the Relationship Between Attachment and Stress, Depression, and Anxiety in Social Work Students. *Clinical Social Work Journal*, 2023;51(4): 339–353. <https://doi.org/10.1007/s10615-023-00882-2>
- Pawlak ND, Serafin L, Czarkowska-Paczek B. Female Sex and Living in a Large City Moderate the Relationships between Nursing Students' Stress Level, Perception of Their Studies, and Intention to Practice Professionally: A Cross-Sectional Study. *International Journal of Environmental Research and Public Health*, 2022;19(9): 5740. <https://doi.org/10.3390/ijerph19095740>
- Wilczek-Ruzyczka E. Empathy as a determinant of perceived stress and styles of coping with stress in medical, law and psychology students. *Acta Neuropsychologica*, 2024;22(2): 151–167. <https://doi.org/10.5604/01.3001.0054.4569>
- Grobecker PA. A sense of belonging and perceived stress among baccalaureate nursing students in clinical placements. *Nurse Education Today*, 2016;36: 178–183. <https://doi.org/10.1016/j.nedt.2015.09.015>
- Lai AY kwan, Lee L, Wang M ping, Feng Y, Lai TT kwan, Ho L ming, et al. Mental Health Impacts of the COVID-19 Pandemic on International University Students, Related Stressors, and Coping Strategies. *Frontiers in Psychiatry*, 2020;11: 584240. <https://doi.org/10.3389/fpsy.2020.584240>
- Clark RM, Besterfield-Sacre M, Dickerson S, Gau D. Impact of COVID-19 on Engineering Students in the Middle Phases of the Pandemic: Academic Motivation, Valued College Experiences, and Stress. *International Journal of Engineering Education*, 2023;39(1): 30–47.
- Brandolin F, Lappalainen P, Gorinelli S, Muotka J, Lappalainen R. Examining the effectiveness and acceptability of a group-based ACT intervention delivered by videoconference to international university students during the COVID-19 pandemic. *Counselling Psychology Quarterly*, 2023; <https://doi.org/10.1080/09515070.2023.2254726>
- Lampe LC, Mueller-Hilke B. Mindfulness-based intervention helps preclinical medical students to contain stress, maintain mindfulness and improve academic success. *BMC Medical Education*, 2021;21(1): 145. <https://doi.org/10.1186/s12909-021-02578-y>

21. Kaya C, Umucu E, Rumrill P, Chan F, Muller V, Bezyak J, et al. Psychometric Validation of a Turkish Version of the Inventory of Common Problems: a Tool for Turkish College Counselors. *International Journal for the Advancement Of Counselling*, 2019;41(3): 436–448. <https://doi.org/10.1007/s10447-018-9360-0>
22. Geng G, Midford R. Investigating First-Year Education Students' Stress Level. *Australian Journal of Teacher EDUCATION*, 2015;40(6): 1–12.
23. Eminoglu DO, Aydin T. Evaluation of Healthy Lifestyle Behaviors and Stress Levels of Dentistry Students during the COVID-19 Pandemic. *Journal of Education and Future-Egitim ve Gelecek Dergisi*, 2022;(22): 85–97. <https://doi.org/10.30786/jef.879958>
24. Zhang Y, Geddes J, Kanga FH, Himelhoch S. Psychological impacts of the COVID-19 pandemic on medical students in the United States of America. *Psychiatry Research*, 2022;313: 114595. <https://doi.org/10.1016/j.psychres.2022.114595>
25. Alkureishi ML, Jaishankar D, Dave S, Tatineni S, Zhu M, Chretien KC, et al. Impact of the Early Phase of the COVID-19 Pandemic on Medical Student Well-Being: a Multisite Survey. *Journal of General Internal Medicine*, 2022;37(9): 2156–2164. <https://doi.org/10.1007/s11606-022-07497-2>
26. Liu B, Huynh E, Li C, Wu Q. Impact of COVID-19 on college students at one of the most diverse campuses in the USA: a factor analysis of survey data. *BMJ Open*, 2022;12(9): e061719. <https://doi.org/10.1136/bmjopen-2022-061719>
27. Martinez L, Robles E, Trofimoff V, Vidal N, Espada AD, Mosquera N, et al. Subjective Well-Being and Mental Health among College Students: Two Datasets for Diagnosis and Program Evaluation. *Data*, 2024;9(3): 44. <https://doi.org/10.3390/data9030044>
28. Pieh C, Dale R, Plener PL, Humer E, Probst T. Stress levels in high-school students after a semester of home-schooling. *European Child & Adolescent Psychiatry*, 2022;31(11): 1847–1849. <https://doi.org/10.1007/s00787-021-01826-2>
29. Marcen-Roman Y, Gasch-Gallen A, Vela Martin de la Mota II, Calatayud E, Gomez-Soria I, Rodriguez-Roca B. Stress Perceived by University Health Sciences Students, 1 Year after COVID-19 Pandemic. *International Journal of Environmental Research and Public Health*, 2021;18(10): 5233. <https://doi.org/10.3390/ijerph18105233>
30. Kostic J, Zikic O, Dordevic V, Krivokapic Z. Perceived stress among university students in south-east Serbia during the COVID-19 outbreak. *Annals of General Psychiatry*, 2021;20(1): 25. <https://doi.org/10.1186/s12991-021-00346-2>
31. Brandolin F, Lappalainen P, Gorinelli AGS, Lappalainen R. Understanding and Explaining Psychological Distress in International Students. *International Journal of Psychology and Psychological Therapy*, 2023;23(1).
32. Tripska K, Draessler J, Pokladnikova J. Heart rate variability, perceived stress and willingness to seek counselling in undergraduate students. *Journal of Psychosomatic Research*, 2022;160: 110972. <https://doi.org/10.1016/j.jpsychores.2022.110972>
33. Leombruni P, Corradi A, Lo Moro G, Acampora A, Agodi A, Celotto D, et al. Stress in Medical Students: PRIMES, an Italian, Multicenter Cross-Sectional Study. *International Journal of Environmental Research and Public Health*, 2022;19(9): 5010. <https://doi.org/10.3390/ijerph19095010>
34. Perello Rank M, Chiro M, Perez de la Ossa P. Stress and burnout in chiropractic students of European chiropractic colleges: A cross-sectional study. *Journal of Chiropractic Education*, 2021;35(1): 14–21. <https://doi.org/10.7899/JCE-19-7>
35. Avunduk ATE, Delikan E. Satisfaction and stress levels of dentistry students relating to distance education. *Dental and Medical Problems*, 2021;58(3): 291–298. <https://doi.org/10.17219/dmp/135318>
36. Toprak ME, Tunc SK. COVID-19 pandemic and Emergency Remote Education Practices: Effects on Dentistry Students. *Nigerian Journal of Clinical Practice*, 2022;25(5): 621–629. https://doi.org/10.4103/njcp.njcp_1564_21
37. Aslan I, Cinar O. Predictors and prevalence of stress, anxiety, depression, and PTSD among university students during the second wave of the COVID-19 pandemic in Turkey. *Frontiers in Psychology*, 2023;13: 1087528. <https://doi.org/10.3389/fpsyg.2022.1087528>
38. Chowdhury U, Suvro MAH, Farhan SMD, Uddin MJ. Depression and stress regarding future career among university students during COVID-19 pandemic. *PLoS One*, 2022;17(4): e0266686. <https://doi.org/10.1371/journal.pone.0266686>
39. Jagoda T, Rathnayake S. Perceived stress and learning environment among nursing students: A cross-sectional study. *Collegian*, 2021;28(5): 587–594. <https://doi.org/10.1016/j.colegn.2021.03.003>
40. Nguyen-Thi TT, Le HM, Chau TL, Le HT, Pham TT, Tran NT, et al. Prevalence of stress and related factors among healthcare students: a cross-sectional study in Can Tho City, Vietnam. *Annali di Igiene Medicina Preventiva e di Comunita*, 2024;36(3): 292–301. <https://doi.org/10.7416/ai.2023.2591>
41. Chen JY, Chin WY, Tiwari A, Wong J, Wong ICK, Worsley A, et al. Validation of the perceived stress scale (PSS-10) in medical and health sciences students in Hong Kong. *Asia Pacific Scholar*, 2021;6(2): 31–37. <https://doi.org/10.29060/TAPS.2021-6-2/OA2328>
42. Alhalaiqa F, Masa'Deh R, Al-Yami M, Al-Ghabeesh S, Rayan A, Shawashreh A, et al. Perceived stress among university students: Syrian refugees versus Jordanians. *Journal of Public Health-Heidelberg*, 2021;29(6): 1477–1486. <https://doi.org/10.1007/s10389-020-01261-8>
43. Wongpakaran N, Oon-Arom A, Karawekpanyawong N, Lohanan T, Leesawat T, Wongpakaran T. Borderline Personality Symptoms: What Not to Be Overlooked When Approaching Suicidal Ideation among University Students. *Healthcare*, 2021;9(10): 1399. <https://doi.org/10.3390/healthcare9101399>

44. Thejaswi SG, Mukerji A, Baliga S, Dewan SK, Verma A. Musculoskeletal pain among medical students and its association with perceived stress level: A cross-sectional study. *Journal of Education and Health Promotion*, 2023;12(1). https://doi.org/10.4103/jehp.jehp_1237_22
45. Sahoo S, Mishra PR, Mishra S, Kar M, Padhy SK. Perceived stress and coping skills in the newly joined medical undergraduate students: An exploratory study from Eastern India. *Journal of Mental Health and Human Behaviour*, 2021;26(2): 122–131. https://doi.org/10.4103/jmhbb.jmhbb_247_21
46. Basheer B, Allassaf M, Alameer A, Alghamdi A, Alzaaqui N, Alfaran F, et al. Emotional intelligence and stress coping mechanisms among dental students in Riyadh, Saudi Arabia. *Medical Science*, 2022;26(119): e2018.
47. Alfreeh L, Abulmeaty MMA, Abudawood M, Aljaser F, Shivappa N, Hebert JR, et al. Association between the Inflammatory Potential of Diet and Stress among Female College Students. *Nutrients*, 2020;12(8): 2389. <https://doi.org/10.3390/nu12082389>
48. Yap SY, Foo CN, Lim YM, Ng FL, Mohd-Sidik S, Tang PY, et al. Traditional Chinese Medicine Body Constitutions and Psychological Determinants of Depression among University Students in Malaysia: A Pilot. *International Journal of Environmental Research and Public Health*, 2021;18(10): 5366. <https://doi.org/10.3390/ijerph18105366>
49. Behzadnia A, Smith DR, Goodson ML. A Cross-sectional Examination of the Relationship between Approaches to Learning and Perceived Stress among Medical Students in Malaysia. *Education for Health*, 2018;31(2): 80–86. https://doi.org/10.4103/efh.EfH_280_17
50. Saddki N, Sukerman N, Mohamad D. Association between Emotional Intelligence and Perceived Stress in Undergraduate Dental Students. *Malaysian Journal of Medical Sciences*, 2017;24(1): 59–68. <https://doi.org/10.21315/mjms2017.24.1.7>
51. Zhan H, Zheng C, Zhang X, Yang M, Zhang L, Jia X. Chinese College Students' Stress and Anxiety Levels Under COVID-19. *Frontiers in Psychiatry*, 2021;12: 615390. <https://doi.org/10.3389/fpsy.2021.615390>
52. Yuan L, Lu L, Wang X, Qu M, Gao Y, Pan B. Comorbid anxiety and depressive symptoms and the related factors among international medical students in China during COVID-19 pandemic: a cross-sectional study. *BMC Psychiatry*, 2023;23(1): 165. <https://doi.org/10.1186/s12888-023-04638-7>
53. Yuan LL, Lu L, Wang XH, Guo XX, Ren H, Gao YQ, et al. Prevalence and Predictors of Anxiety and Depressive Symptoms Among International Medical Students in China During COVID-19 Pandemic. *Frontiers in Psychiatry*, 2021;12: 761964. <https://doi.org/10.3389/fpsy.2021.761964>
54. Jia N, Sakulsriprasert C, Wongpakaran N, Suradom C, O' Donnell R. Borderline Personality Disorder Symptoms and Its Clinical Correlates among Chinese University Students: A Cross-Sectional Study. *Healthcare*, 2022;10(9): 1751. <https://doi.org/10.3390/healthcare10091751>
55. Ma Y, Zhang B, Meng Y, Cao Y, Mao Y, Qiu C. Perceived stress and depressive symptoms among Chinese college students: A moderated mediation model of biorhythm and ego resilience. *Frontiers in Public Health*, 2022;10: 951717. <https://doi.org/10.3389/fpubh.2022.951717>
56. Lin Y, Zhang X, Li H, Huang Y, Zhang W, Zhang C. Musculoskeletal pain is prevalent in Chinese medical and dental students: A cross-sectional study. *Frontiers in Public Health*, 2022;10: 1046466. <https://doi.org/10.3389/fpubh.2022.1046466>
57. Qiao N. Does Perceived Stress of University Students Affected by Preferences for Movie Genres? an Exploratory Cross-Sectional Study in China. *Frontiers in Psychology*, 2021;12: 761340. <https://doi.org/10.3389/fpsyg.2021.761340>
58. Zhang Y, Jia Y, MuLaTiHaJi M, Mi Y, Mei Y, Sun T, et al. A cross-sectional mental-health survey of Chinese postgraduate students majoring in stomatology post COVID-19 restrictions. *Frontiers in Public Health*, 2024;12: 1376540. <https://doi.org/10.3389/fpubh.2024.1376540>
59. Ochnik D, Rogowska AM, Kusnierz C, Jakubiak M, Schuetz A, Held MJ, et al. Mental health prevalence and predictors among university students in nine countries during the COVID-19 pandemic: a cross-national study. *Scientific Reports*, 2021;11(1): 18644. <https://doi.org/10.1038/s41598-021-97697-3>
60. Mocny-Pachonska K, Trzcionka A, Doniec RJ, Siecinski S, Tanasiewicz M. The Influence of Gender and Year of Study on Stress Levels and Coping Strategies among Polish Dental. *Medicina-Lithuania*, 2020;56(10): 531. <https://doi.org/10.3390/medicina56100531>
61. Wroblewski O, Michalczyk K, Kozlowski M, Nowak K, Chudecka-Glaz A, Skwirczynska E. A Longitudinal Study of the Influence of the COVID-19 Pandemic on Anxiety and Stress among Medical University Students. *Journal of Clinical Medicine*, 2024;13(3): 890. <https://doi.org/10.3390/jcm13030890>
62. Badura-Brzoza K, Debski P, Glowczynski P, Debska-Janus M, Gorczyca P. Life Satisfaction and Perceived Stress versus Health Promoting Behavior among Medical Students during the COVID-19 Pandemic. *International Journal of Environmental Research and Public Health*, 2022;19(11): 6706. <https://doi.org/10.3390/ijerph19116706>
63. Kupcewicz E, Grochans E, Kaducakova H, Mikla M, Jozwik M. Analysis of the Relationship between Stress Intensity and Coping Strategy and the Quality of Life of Nursing Students in Poland, Spain and Slovakia. *International Journal of Environmental Research and Public Health*, 2020;17(12): 4536. <https://doi.org/10.3390/ijerph17124536>
64. Zdun-Ryzewska A, Sobczak K, Rudnik A. Fatigue, Pro-Social Attitude and Quality of Life as Predictors of Empathy in Medical and Social-Oriented Students. *International Journal of Environmental Research and Public Health*, 2022;19(23): 15853. <https://doi.org/10.3390/ijerph192315853>
65. Shpakou A, Krajewska-Kulak E, Cybulski M, Sokolowska D, Andryszczyk M, Kleszczewska E, et

- al. Anxiety, Stress Perception, and Coping Strategies among Students with COVID-19 Exposure. *Journal of Clinical Medicine*, 2023;12(13): 4404. <https://doi.org/10.3390/jcm12134404>
66. Marchewka W, Kluczevska M, Ciszek K, Zygmunt M, Popiolek L, Marchewka J, et al. Assessment of stress load and its causes among dental students. *Psychiatria Polska*, 2022;56(5): 979–990. <https://doi.org/10.12740/PP/140368>
67. Shpakou A, Krajewska-Kulak E, Cybulski M, Seredocha I, Talaj A, Andryszczyk M, et al. Stress Perception and Coping Strategies of Students on Both Sides of the EU's Eastern Border during the COVID-19 Pandemic. *International Journal of Environmental Research and Public Health*, 2022;19(16): 10275. <https://doi.org/10.3390/ijerph191610275>
68. Balgiu BA, Sfeatcu R, Dumitrache MA, Tribus L. Construct validity and reliability of perceived stress scale (pss-14) in a romanian sample of dental students. *International Journal of Medical Dentistry*, 2021;25(2): 23–30.
69. Butnaru GI, Haller AP, Dragolea LL, Anichiti A, Harsan GDT. Students' Wellbeing during Transition from Onsite to Online Education: Are There Risks Arising from Social Isolation? *International Journal of Environmental Research and Public Health*, 2021;18(18): 9665. <https://doi.org/10.3390/ijerph18189665>
70. Ionescu CG, Chendea A, Licu M. Is Satisfaction with Online Learning Related to Depression, Anxiety, and Insomnia Symptoms? A Cross-Sectional Study on Medical Undergraduates in Romania. *European Journal of Investigation in Health Psychology and Education*, 2023;13(3): 580–594. <https://doi.org/10.3390/ejihpe13030045>
71. Palos R. Students' core self-evaluations and academic burnout: the mediating role of personal resources. *Journal of Applied Research in Higher Education*, 2024;16(3): 904–918. <https://doi.org/10.1108/JARHE-04-2023-0152>
72. Puiu S, Idowu SO, Meghisan-Toma GM, Badircea RM, Doran NM, Manta AG. Online Education Management: A Multivariate Analysis of Students' Perspectives and Challenges during Online Classes. *Electronics*, 2023;12(2): 454. <https://doi.org/10.3390/electronics12020454>
73. Simionescu M, Pellegrini A, Bordea EN. The Effects of COVID-19 Pandemic on Stress Vulnerability of Nursing Students According to Labour Market Status. *Healthcare*, 2021;9(6): 633. <https://doi.org/10.3390/healthcare9060633>
74. Vintila M, Tudorel OI, Stefanut A, Ivanoff A, Bucur V. Emotional distress and coping strategies in COVID-19 anxiety. *Current Psychology*, 2023;42(20): 17503–17512. <https://doi.org/10.1007/s12144-021-02690-8>
75. Dumitrache L, Stanculescu E, Nae M, Dumbraveanu D, Simion G, Talos AM, et al. Post-Lockdown Effects on Students' Mental Health in Romania: Perceived Stress, Missing Daily Social Interactions, and Boredom Proneness. *International Journal of Environmental Research and Public Health*, 2021;18(16): 8599. <https://doi.org/10.3390/ijerph18168599>
76. Lovin D, Bernardeau-Moreau D. Stress among Students and Difficulty with Time Management: A Study at the University of Galati in Romania. *Social Sciences-Basel*, 2022;11(12): 538. <https://doi.org/10.3390/socsci11120538>
77. Vlad R, Golu F, Toma A, Draganescu D, Oprea B, Chiper BI. Depression and anxiety in romanian medical students: prevalence and associations with personality. *Farmacia*, 2020;68(5): 944–949. <https://doi.org/10.31925/farmacia.2020.5.24>
78. Jansen N, Kuehl J, Eller N, Kingdon P, Firk C. Mentalizing Capacity Partially Mediates the Relationship Between Attachment and Stress, Depression, and Anxiety in Social Work Students. *Clinical Social Work Journal*, 2023;51(4): 339–353. <https://doi.org/10.1007/s10615-023-00882-2>
79. Iermakov S, Yermakova T, Prusik K. Modeling the gameplay actions of elite volleyball players and teams based on statistical match reports. *Pedagogy of Health*. 2023;2(2):50–64. <https://doi.org/10.15561/health.2023.0202>
80. Yermakova T, Lutskyi V, Piatnytska D. Motivation for physical activity among university students residing in areas of military conflict in Ukraine. *Pedagogy of Health*. 2023;2(2):74–82. <https://doi.org/10.15561/health.2023.0204>
81. Kryventsova I, Potop V, Prusik K, Samoilo D. Development of coordination abilities of fencers aged 10–12 years living in the territory of martial law. *Pedagogy of Health*. 2022;1(2):34–8. <https://doi.org/10.15561/health.2022.0201>
82. Golenkova J, Kravchuk T, Sanzharova N, Potop V, Filon K. Psychological recreation of overcoming failures and achieving success by young rhythmic gymnasts aged 6–8. *Physical Culture, Recreation and Rehabilitation*. 2023;2(1):4–9. <https://doi.org/10.15561/physcult.2023.0101>
83. Vypasniak I, Shesterova L, Kryventsova I. The professional readiness of student-teachers in physical education in Ukraine's war-torn areas. *Pedagogy of Health*. 2023;2(2):83–90. <https://doi.org/10.15561/health.2023.0205>
84. Ciavoi G, Marginean I, Tig IA, Todor L, Farcasiu AT, Farcasiu C, et al. Test anxiety in dental medical students during covid-19 pandemic. *Romanian Journal of Oral Rehabilitation*, 2021;13(2): 52–59.
85. Stefenel D, Gonzalez JM, Rogobete S, Sassu R. Coping Strategies and Life Satisfaction among Romanian Emerging Adults during the COVID-19 Pandemic. *Sustainability*, 2022;14(5): 2783. <https://doi.org/10.3390/su14052783>

Information about the authors:

Vladimir Potop; <https://orcid.org/0000-0001-8571-2469>; vladimir_potop@yahoo.com; Doctoral School of Sports Science and Physical Education, National University of Science and Technology Polytechnic Bucharest, University Center Pitești (Pitesti, Romania); Department of Physical Education and Sport, National University of Science and Technology Polytechnic Bucharest, University Center Pitești (Pitesti, Romania); State University of Physical Education and Sport (Chisinau, Republic of Moldova).

Igor Vypasniak; (Corresponding Author); <https://orcid.org/0000-0002-4192-1880>; ihor.vypasniak@pnu.edu.ua; Department of Theory and Methods of Physical Culture, Vasyl Stefanyk Precarpathian National University; Ivano-Frankivsk, Ukraine.

Iryna Ivanyshyn; <https://orcid.org/0000-0003-1765-8311>; iryna.ivanyshyn@pnu.edu.ua; Department of Theory and Methods of Physical Culture, Vasyl Stefanyk Precarpathian National University; Ivano-Frankivsk, Ukraine.

Vasyl Lutskiy; <https://orcid.org/0000-0003-3940-1349>; luckij55@gmail.com; Department of Theory and Methods of Physical Culture; Vasyl Stefanyk Precarpathian National University; Ivano-Frankivsk, Ukraine.

Iryna Kryventsova; PhD of Pedagogical Sciences, Associate Professor; <https://orcid.org/0000-0001-6931-3978>; kriventsova.ira@ukr.net; Department of Theory, Methodology and Practice of Physical Education; H. S. Skovoroda Kharkiv National Pedagogical University; Kharkiv, Ukraine.

Lyudmila Shesterova; PhD (Physical Education and Sport), Professor; The Head of the Department; <https://orcid.org/0000-0001-8777-6386>; lydmula121056@gmail.com; Department of Theory and Methodology of Physical Education; Municipal Establishment “Kharkiv Humanitarian-Pedagogical Academy” of the Kharkiv Regional Council; 7, Rustaveli Lane, 61001, Kharkiv, Ukraine.

Katarzyna Prusik; <https://orcid.org/0000-0002-2960-5105>; prusikkatarzyna@gmail.com; Department of Sports, Gdansk University of Physical Education and Sport; Gdansk, Poland.

Cite this article as:

Potop V, Vypasniak I, Ivanyshyn I, Lutskiy V, Kryventsova I, Shesterova L, Prusik K. Assessment of stress and health conditions among students in the context of the war in Ukraine. *Physical Culture, Recreation and Rehabilitation*, 2024;3(2):58–69.

<https://doi.org/10.15561/physcult.2024.0203>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited (<http://creativecommons.org/licenses/by/4.0/deed.en>).

Received: 22.08.2024

Accepted: 11.10.2024; Published: 30.12.2024